

DISCRETE LINEAR GROUPS CONTAINING ARITHMETIC GROUPS

INDIRA CHATTERJI AND T. N. VENKATARAMANA

ABSTRACT. If H is a simple real algebraic subgroup of real rank at least two in a simple real algebraic group G , we prove, in a substantial number of cases, that a Zariski dense discrete subgroup of G containing a lattice in H is a lattice in G . For example, we show that any Zariski dense discrete subgroup of $SL_n(\mathbb{R})$ ($n \geq 4$) which contains $SL_3(\mathbb{Z})$ (in the top left hand corner) is commensurable with a conjugate of $SL_n(\mathbb{Z})$.

In contrast, when the groups G and H are of real rank one, we prove that a lattice Δ in a real rank one group H embedded in a larger real rank one group G extends to a Zariski dense discrete subgroup Γ of G of **infinite co-volume**.

1. INTRODUCTION

In this paper, we study special cases of the following problem raised by Madhav Nori:

Problem 1. (*Nori, 1983*) *If H is a real algebraic subgroup of a real semi-simple algebraic group G , find sufficient conditions on H and G such that any Zariski dense subgroup Γ of G which intersects H in a lattice in H , is itself a lattice in G .*

If the smaller group H is a simple Lie group and has real rank strictly greater than one (then the larger group G is also of real rank at least two), we do not know any example when the larger discrete group Γ is not a lattice. The goal of the present paper is to study the following question, related to Nori's problem.

Question 2. *If a Zariski dense discrete subgroup of a simple non-compact Lie group G intersects a simple non-compact Lie subgroup H*

1991 *Mathematics Subject Classification.* Primary 22E40, Secondary 20G30
Indira Chatterji, Université d'Orléans, Route de Chartres, 45000 Orléans, FRANCE. Partially supported by NSF grant DMS 0644613.
T. N. Venkataramana, School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road, Bombay - 400 005, INDIA. Supported by J.C. Bose Fellowship for the period 2008-2013.

in a lattice, is the larger discrete group a lattice in the larger Lie group G ?

Analogous questions have been considered before: (e.g. [F-K], Bass-Lubotzky [B-L], Hee Oh [O]), and [V2]). In the present paper, we give some positive evidence for this conjecture.

For example, consider the “top left hand corner” embedding

$$SL_k \subset SL_n, \quad (k \geq 3).$$

The embedding is as follows: an SL_k matrix M is thought of as an $n \times n$ matrix M' such that the first $k \times k$ entries of M' are the same as those of M , the last $(n - k) \times (n - k)$ entries of M' are those of the identity $(n - k) \times (n - k)$ matrix, and all other entries of M' are zero.

Theorem 3. *Suppose that SL_3 is embedded in SL_n (in the “top left hand corner”) as above. Suppose that Γ is a Zariski dense discrete subgroup of $SL_n(\mathbb{R})$ whose intersection with $SL_3(\mathbb{R})$ is a subgroup of $SL_3(\mathbb{Z})$ of finite index. Then, Γ is commensurate to a conjugate of $SL_n(\mathbb{Z})$, and is hence a lattice in $SL_n(\mathbb{R})$.*

We may similarly embed $Sp_k \subset Sp_g$ in the “top left hand corner”, where Sp_g is the symplectic group of $2g \times 2g$ -matrices preserving a non-degenerate symplectic form in $2g$ -variables. Denote by $Sp_g(\mathbb{Z})$ the integral symplectic group.

Theorem 4. *If Γ is a Zariski dense discrete subgroup of $Sp_g(\mathbb{R})$ whose intersection with $Sp_2(\mathbb{R})$ is commensurate to $Sp_2(\mathbb{Z})$ then a conjugate of Γ is commensurate with $Sp_g(\mathbb{Z})$ and hence Γ is a lattice in $Sp_g(\mathbb{R})$.*

The proof of Theorem 3 (as does the proof of Theorem 4) depends on a general super-rigidity theorem for discrete subgroups Γ which contain a “large” enough higher rank lattice. More precisely, our main result is the following.

Let P be a minimal real parabolic subgroup of the simple Lie group G and let denote by S and N respectively a maximal real split torus of P and the unipotent radical of P . Let A be the connected component of identity in S . Denote by P_0 the subgroup AN of P (if G is split, then $P_0 = P$; in general we have replaced the minimal parabolic subgroup P by a subgroup P_0 which has no compact factors such that P/P_0 is compact). Let K be a maximal compact subgroup of G . We have the Iwasawa decomposition $G = P_0K = ANK$. Assume that the isotropies of H in its action on the quotient G/P_0 are all positive dimensional and non-compact.

Theorem 5. *Let H be a semi-simple subgroup of the simple Lie group G with $\mathbb{R}\text{-rank}(H) \geq 2$. Let Γ be a Zariski dense subgroup of G whose intersection with H is an irreducible lattice in H . Let $G = NAK$ be the Iwasawa decomposition of G as above and $P_0 = AN$. If the isotropy of H acting on G/P_0 is positive dimensional at every point of G/P_0 , then Γ is a super-rigid subgroup of G .*

The conditions of Theorem 5 are satisfied if the dimension of H is sufficiently large (for example, if $\dim(K) < \dim(H)$). We then get the following Theorem as a corollary of Theorem 5.

Theorem 6. *Let Γ be a Zariski dense discrete subgroup of a simple Lie group G which intersects a semi-simple subgroup H of G (with $\mathbb{R}\text{-rank}(H) \geq 2$) in an irreducible lattice. Let K be a maximal compact subgroup of G and assume that $\dim(H) > \dim(K)$. Then Γ is a super-rigid subgroup of G .*

Here, super-rigid is in the sense of Margulis [M]. That is, all linear representations - satisfying some mild conditions - of the group Γ virtually extend to (i.e. coincide on a finite index subgroup of Γ with) a linear representation of the ambient group G . If Nori's question is in the affirmative, then Γ must be a lattice and by the Super-rigidity Theorem of Margulis, must satisfy the super-rigidity property; Theorem 6 is therefore evidence that the answer to our question is "yes".

It was Madhav Nori who first raised the question (to the second named author of the present paper) whether these larger discrete groups containing higher rank lattices have to be lattices themselves (i.e. arithmetic groups, in view of Margulis' Arithmeticity Theorem). We will in fact show several examples of pairs of groups (H, G) , such that Γ does turn out to be a lattice, confirming Nori's guess (see Theorem 3 and Corollary 1).

For instance, as a consequence of Theorem 6 we obtain the following.

Corollary 1. *If $n \geq 4$ and $H = SL_{n-1}(\mathbb{R}) \subset SL_n(\mathbb{R})$ and Γ is a Zariski dense subgroup of $SL_n(\mathbb{R})$ which intersects $SL_{n-1}(\mathbb{R})$ in a finite index subgroup of $SL_{n-1}(\mathbb{Z})$, then a conjugate of Γ in $SL_n(\mathbb{R})$ is commensurate to $SL_n(\mathbb{Z})$.*

Analogously, we prove

Corollary 2. *If $g \geq 3$, then under the standard embedding $Sp_{g-1} \subset Sp_g$, every Zariski dense discrete subgroup of $Sp_g(\mathbb{R})$ which contains a*

finite index subgroup of $Sp_{g-1}(\mathbb{Z})$ is commensurable to a conjugate of $Sp_g(\mathbb{Z})$.

The proof of Theorem 5 runs as follows.

We adapt Margulis' proof of super-rigidity to our situation; the proof of Margulis uses crucially that a lattice Γ acts ergodically on G/S for any non-compact subgroup S of G , whereas we do not have the ergodicity available to us. We use instead the fact that the representation of the discrete group Γ is rational on the smaller group Δ .

Given a representation ρ of the group Γ on a vector space over a local field k' , we use a construction of Furstenberg to obtain a Γ -equivariant measurable map ϕ from G/P_0 into the space $\mathcal{P}$ of probability measures on the projective space of the vector space. Using the fact that the isotropy subgroup of H at any point in G/P_0 is non-compact, we deduce that on H -orbits, the map ϕ is rational, and by pasting together the rationality of many such orbits, we deduce the rationality of the representation ρ .

Theorem 6 implies Corollary 1 as follows: the super-rigidity of Γ in Corollary 1 implies, as in [M], that Γ is a *subgroup* of an arithmetic subgroup Γ_0 of $SL_n(\mathbb{R})$. It follows that the $\mathbb{Q}$ -form of $SL_n(\mathbb{R})$ associated to this arithmetic group has $\mathbb{Q}$ -rank greater than $n/2$. The classification of the $\mathbb{Q}$ -forms of $SL_n(\mathbb{R})$ then implies that the $\mathbb{Q}$ -form must be $SL_n(\mathbb{Q})$ and that Γ is commensurate to a subgroup of $SL_n(\mathbb{Z})$. Since Γ is Zariski dense and virtually contains $SL_{n-1}(\mathbb{Z})$, it follows from [V1] that Γ is commensurate to $SL_n(\mathbb{Z})$.

We now give a list (not exhaustive) of more pairs (H, G) which satisfy the condition $(\dim(H) > \dim(K))$ of Theorem 6.

Corollary 3. *If (H, G) is one of the pairs*

- (1) $H = Sp_g \subset SL_{2g}$ with $g \geq 2$, or
- (2) $H = SL_p \times SL_p \subset G = SL_{2p}$ with $p \geq 3$
- (3) $H = Sp_a \times Sp_a \subset G = Sp_{2a}$ with $1 \leq a$.

then any Zariski dense discrete subgroup Γ of $G(\mathbb{R})$ whose intersection with $H(\mathbb{R})$ is an irreducible lattice, is super-rigid in $G(\mathbb{R})$.

There are examples of pairs (H, G) satisfying the condition of Theorem 5 which are not covered by Theorem 6. In the cases of the following Corollary, it is easy to check that $\dim(H) = \dim(K)$ and that the isotropy of H at any generic point of G/P is a non-compact Cartan subgroup of H . When $G = H(\mathbb{C})$, we view G as the group of **real**

points of a complex algebraic group, and Zariski density of a subgroup $\Gamma \subset G$ is taken to mean that Γ is Zariski dense in $G(\mathbb{C}) = H(\mathbb{C}) \times H(\mathbb{C})$.

Corollary 4. *Let H be a real simple Lie group with $\mathbb{R} - \text{rank}(H) \geq 2$ embedded in the complex group $G = H(\mathbb{C})$. If H has no compact Cartan subgroup, then every Zariski dense discrete subgroup Γ of G which intersects H in a lattice is super-rigid in G .*

Notice that Theorem 6 and Theorem 5 allow us to deduce that if Γ is as in Theorem 6 or Theorem 5, then Γ 's is a subgroup of an arithmetic group in G (see Theorem 20), reducing Nori's question to the following apparently simpler one:

Question 7. *If a Zariski dense subgroup Γ of a lattice in a simple non-compact Lie group G contains a higher rank lattice of a smaller group, is Γ itself a lattice in G ?*

We now describe the proof of Theorem 3. If Theorem 6 is to be applied directly, then the dimension of the maximal compact of $SL_n(\mathbb{R})$ must be less than the dimension of $SL_3(\mathbb{R})$, which can only happen if $n = 4$; instead, what we will do, is to show that the group generated by $SL_3(\mathbb{Z})$ (in the top left hand corner of $SL_n(\mathbb{R})$ as in the statement of Theorem 3) and a conjugate of a unipotent root subgroup of $SL_3(\mathbb{Z})$ by a generic element of Γ , (modulo its radical), is a Zariski dense *discrete* subgroup of $SL_4(\mathbb{R})$. By applying Corollary 1 for the pair $SL_3(\mathbb{R})$ and $SL_4(\mathbb{R})$, we see that the Zariski dense discrete subgroup Γ of $SL_n(\mathbb{R})$ contains, virtually, a conjugate of $SL_4(\mathbb{Z})$.

We can apply the same procedure to SL_4 instead of SL_3 and obtain $SL_5(\mathbb{Z})$ as a subgroup of Γ , ..., and finally obtain that Γ virtually contains a conjugate of $SL_n(\mathbb{Z})$. This proves Theorem 3. The proof of Theorem 4 is similar: use Corollary 2 in place of Corollary 1.

When G has real rank one, we show (Theorem 24), that the answer to the counterpart of Question 2 is in the negative! That is, given $H \subset G$ with $\mathbb{R} - \text{rank}(G) = 1$, and a lattice $\Delta \subset H$, one can always produce a Zariski dense discrete subgroup Γ in G which is not a lattice in G , whose intersection with H is a subgroup of finite index in Δ . The method is essentially that of Fricke and Klein [F-K] who produce, starting from Fuchsian groups, Kleinian groups of infinite co-volume, by using a "ping-pong" argument.

2. PRELIMINARIES ON MEASURABLE MAPS

The aim of this section is to recall a few well known facts used in the proof of Theorem 6 and prove the following key fact.

Proposition 8. *Let $\tilde{H}$ be a $\mathbb{Q}$ -simple, simply connected algebraic group with $\mathbb{R}$ -rank $(H) \geq 2$ and $\Delta \subset \tilde{H}(\mathbb{Z})$ an arithmetic subgroup. Let $\rho : \Delta \rightarrow G'(k')$ be a representation, where G' is a linear algebraic group over a local field k' of characteristic zero. Suppose that $S \subset \tilde{H}(\mathbb{R})$ a closed non-compact subgroup and let $J \subset G'(k')$ be an algebraic subgroup. Let $\phi : \tilde{H}(\mathbb{R})/S \rightarrow G'(k')/J$ be a Borel measurable map which is Δ -equivariant. Then the map $\phi : \tilde{H}(\mathbb{R})/S \rightarrow G'(k')/J$ is a rational map. More precisely:*

If k' is an archimedean local field, there exist a homomorphism $\tilde{\rho} : \tilde{H}(\mathbb{R}) \rightarrow G'(k')$ of real algebraic groups defined over $\mathbb{R}$ and a point $p \in G'(k')/J$, such that the map $\phi(h) = \tilde{\rho}(h)(p)$ for all $h \in \tilde{H}(\mathbb{R})$ (and is therefore an $\mathbb{R}$ -rational map of real varieties).

If k' is a non-archimedean local field, then the map ϕ is constant.

An important ingredient in the proof is the following generalisation of the Ergodicity Theorem of Moore (see [Z], Theorem (2.2.6)).

Lemma 9. *Let $\tilde{H}$ be a $\mathbb{Q}$ -simple, simply connected algebraic group with $\mathbb{R}$ -rank $(H) \geq 1$ and $\Delta \subset \tilde{H}(\mathbb{Z})$ an arithmetic subgroup. For any closed non-compact subgroup $S \subset \tilde{H}(\mathbb{R})$, the group Δ acts ergodically on the quotient $\tilde{H}(\mathbb{R})/S$.*

The $\mathbb{Q}$ -simplicity of $\tilde{H}$ implies that there exists a connected absolutely almost simple simply connected group H_0 defined over a number field K , such that $\tilde{H} = R_{K/\mathbb{Q}}(H_0)$, where $R_{K/\mathbb{Q}}(H_0)$ is the Weil restriction of H_0 from K to $\mathbb{Q}$. Consequently,

$$\tilde{H}(\mathbb{R}) = H_0(K \otimes \mathbb{R}) = \prod_{\alpha \in \infty} H_0(K_\alpha),$$

where ∞ denote the set of equivalence classes of archimedean embeddings of K . Let $A \subset \infty$ denote the archimedean embeddings $\{\alpha\}$ such that $H_0(K_\alpha)$ is non-compact. Then, $\tilde{H} = H_u \times H^*$ where $H_u := \prod_{\alpha \in \infty \setminus A} H_0(K_\alpha)$ is a compact group, and $H^* := \prod_{\alpha \in A} H_0(K_\alpha)$ is a semi-simple group without compact factors.

Lemma 10. *Let $\tilde{H}$ be a $\mathbb{Q}$ -simple, simply connected algebraic group with $\mathbb{R}$ -rank $(H) \geq 1$ and $\Delta \subset \tilde{H}(\mathbb{Z})$ an arithmetic subgroup. Suppose*

that π is a unitary representation of $\tilde{H}(\mathbb{R})$ on a Hilbert space, such that for any simple factor H_α of H^* , the space π^{H_α} of vectors of π invariant under the subgroup H_α is zero. Then, the space π^S of vectors in π invariant under the non-compact subgroup $S \subset \tilde{H}(\mathbb{R})$ is also zero.

Proof. By the Howe-Moore theorem, (see [Z], Theorem (2.2.20)), there exists a proper function $\Xi : H^* \rightarrow [0, +\infty)$ (where $[0, +\infty)$ is the closed open interval with end points 0 and $+\infty$) with the following property. Let π be a unitary representation of H^* on a Hilbert space V such that for any non-compact simple factor $H_\alpha := H_0(K_\alpha)$ the space of invariants V^{H_α} is zero. For $v, w \in V$, denote their inner product by $\langle v, w \rangle$ and define $|v|$ by the formula $|v|^2 = \langle v, v \rangle$. Then, for all $v, w \in V$, and all $g \in H^*$ we have

$$|\langle \pi(g)v, w \rangle| \leq \frac{|v| |w|}{\Xi(g)}.$$

Suppose now that π is a unitary representation of $\tilde{H}(\mathbb{R})$ on a Hilbert space such that for any non-compact simple factor H_α of H^* , the space of H_α invariants in π is zero. Let $g \in S \subset \tilde{H}(\mathbb{R}) = H^* \times H_u$; we write $g = (g^*, g_u)$ accordingly. Then, by the estimate of the preceding paragraph applied to $g^* \in H^*$ and the vectors $\pi(g_u)v$ in place of v , and v in place of w , we get:

$$|\langle \pi(g)v, v \rangle| \leq \frac{|v| |v|}{\Xi(g^*)}.$$

If π has non-zero S -invariant vectors v (say, of norm 1), then the foregoing estimate shows that $\Xi(g^*)$ is bounded on S , which by the properness of Ξ implies that S is compact; since this is false by assumption, we have proved that π does not have any non-zero vectors invariant under S . $\square$

We are now ready to complete the proof of Lemma 9.

Proof of Lemma 9. Let $V_0 = L^2[\Delta \backslash \tilde{H}(\mathbb{R})]$ be the space of square integrable functions on $\Delta \backslash \tilde{H}(\mathbb{R})$; the latter space has finite volume, and hence contains the space of constant functions. For any simple factor H_α of $H^* \subset \tilde{H}(\mathbb{R})$, the space $V_0^{H_\alpha}$ is just the space of constants, by strong approximation (see [M], Chapter (II), Theorem (6.7); in the notation of [M], we may take $B = \{\alpha\}$ to be a singleton). Consequently, if π denotes the space of functions in V_0 orthogonal to the constant functions, then π satisfies the assumptions of Lemma 10. Therefore, by Lemma 10, $\pi^S = 0$, and hence the only functions on $\Delta \backslash \tilde{H}(\mathbb{R})$ invariant under the non-compact group S are constants. This proves Lemma 9. $\square$

We now record a consequence of Lemma 9 which will be used in the proof of Proposition 8.

Lemma 11. *Let $\tilde{H}$ be a $\mathbb{Q}$ -simple, simply connected algebraic group with $\mathbb{R}\text{-rank}(\tilde{H}) \geq 1$ and $\Delta \subset \tilde{H}(\mathbb{Z})$ an arithmetic subgroup. Suppose that $s \in \tilde{H}(\mathbb{R})$ generates an infinite discrete subgroup and let $\tau : s^{\mathbb{Z}} \rightarrow \mathbb{Z}$ be an isomorphism. Then, any $s^{\mathbb{Z}}$ -equivariant Borel measurable map*

$$\phi^* : \Delta \backslash \tilde{H}(\mathbb{R}) \rightarrow \mathbb{Z}$$

is constant.

Proof. The group $s^{\mathbb{Z}}$ generated by s is non-compact, therefore according to Lemma 9, for every $m \neq 0$, the group $s^{m\mathbb{Z}}$ acts ergodically on $\Delta \backslash \tilde{H}(\mathbb{R})$. Since $\mathbb{Z}$ is discrete, the orbits of $s^{m\mathbb{Z}}$ in $\mathbb{Z}$ are closed. Composing ϕ^* with the projection $\mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$ gives an $s^{m\mathbb{Z}}$ -equivariant map

$$\overline{\phi^*} : \Delta \backslash \tilde{H}(\mathbb{R}) \rightarrow \mathbb{Z}/m\mathbb{Z}.$$

Hence by Proposition (2.1.11) of [Z], the map $\overline{\phi^*}$ is essentially constant so that the image of ϕ^* is essentially contained in an orbit of $s^{m\mathbb{Z}}$; that is, there exists a set E_m of measure zero in $\Delta \backslash \tilde{H}(\mathbb{R})$ such that the image under ϕ^* of the complement F_m of E_m is contained in an orbit of $s^{m\mathbb{Z}}$. The intersection of all these orbits is non-empty, since it contains the image of the countable intersection $\cap_{m \neq 0} F_m$, which is a set of full measure. In that case, the image must be a point: if p, q lie in the image, then for every $m \neq 0$, there exists an integer n such that $q = \tau(s^{mn})(p)$; that is $p - q = mn\tau(s) = \pm mn\tau(s)$; therefore $p - q = 0$. $\square$

The following is a version of the Margulis super-rigidity theorem, except that the Zariski closure of the image $\rho(\Delta)$ is not assumed to be an absolutely simple group and that $\rho(\Delta)$ is not assumed to have non-compact closure in $G'(k')$ if k' is archimedean.

Theorem 12. (Margulis) *Let $\tilde{H}$ be a $\mathbb{Q}$ -simple simply connected algebraic group defined over $\mathbb{Q}$ of $\mathbb{R}\text{-rank}(\tilde{H}) \geq 2$, $\Delta \subset \tilde{H}(\mathbb{Z})$ an arithmetic subgroup and $\rho : \Delta \rightarrow G'(k')$ a homomorphism into a linear algebraic group over a local field k' of characteristic zero.*

(1) ([M], Chapter(VIII), Theorem (3.4) part(a)): *If k' is archimedean, then the map ρ coincides, on a subgroup of finite index, with a representation $\tilde{\rho} : \tilde{H}(\mathbb{R}) \rightarrow G'(k')$.*

(2) ([M], Chapter(VIII), Theorem (5.9) part(a)): *If the local field k' is non-archimedean, then $\rho(\Delta)$ is contained in a compact subgroup of $G'(k')$.*

We can now prove the main result (Proposition 8) of this section.

Proof of Proposition 8. Suppose first that k' is archimedean. Let $\Delta' \subset \Delta$ be a subgroup of finite index such that there exists (according to Theorem 12 quoted above) a representation $\tilde{\rho} : \tilde{H}(\mathbb{R}) \rightarrow G'(k')$ which coincides with ρ on Δ' . Consider the map $\phi^*(h) = \tilde{\rho}(h)^{-1}(\phi(h))$ from $\tilde{H}(\mathbb{R})$ into the quotient $G'(k')/J$. Then, for all $\delta \in \Delta'$, $h \in \tilde{H}(\mathbb{R})$, $s \in S$, we have $\phi^*(\delta h) = \phi^*(h)$ and $\phi^*(hs) = \tilde{\rho}(s)^{-1}(\phi^*(h))$. That is, the map ϕ^* is Δ' invariant and S -equivariant for the action of $\tilde{H}(\mathbb{R})$ on $G'(k')/J$ via the representation $\tilde{\rho}$.

The representation $\tilde{\rho}$ is algebraic; moreover, since k' is archimedean, by assumption the group J is a real algebraic subgroup of $G'(k')$, and hence the action of $\tilde{H}(\mathbb{R})$ on $G'(k')/J$ is smooth. Let S_1 denote the Zariski closure of the image $\tilde{\rho}(S)$. The S_1 -action on $G'(k')/J$ is smooth, hence the quotient $S_1 \backslash G'(k')/J$ is countably separated. On the other hand, by Lemma 9, the action of S on $\Delta' \backslash \tilde{H}(\mathbb{R})$ is ergodic. Hence, by Proposition (2.1.11) of [Z], the image of ϕ^* is essentially contained in an S_1 -orbit i.e. there exists a Borel set E of measure zero in $\tilde{H}(\mathbb{R})/S$, such that the image under ϕ of the complement of E is contained in an S_1 -orbit.

Since S is a non-compact Lie group, S contains an element s of infinite order which generates a discrete non-compact subgroup. We may replace S by the closure of the group generated by the element s and assume that S is abelian. Similarly, we may replace S_1 by the Zariski closure of the image of $s^{\mathbb{Z}}$ and assume that S_1 is also abelian. Hence the S_1 -orbit of the preceding paragraph is of the form S_1/S_2 with S_2 an algebraic subgroup of the abelian group S_1 .

Suppose that the inverse image $S' = S \cap \tilde{\rho}^{-1}(S_2)$ is a non-compact subgroup. By Lemma 9, the group S' acts ergodically on $\Delta \backslash \tilde{H}(\mathbb{R})$; therefore, the map ϕ^* - being S' invariant - is constant: $\phi^*(\Delta \backslash \tilde{H}(\mathbb{R})) = \{p\}$ for some point $p \in G'(k')/J$. That is $\phi(h) = \tilde{\rho}(h)(p)$, and is rational.

Suppose that S' is compact. Since $s^{\mathbb{Z}}$ generates a discrete non-compact subgroup and S' is compact, the image $s_1^{\mathbb{Z}} := \tilde{\rho}(s^{\mathbb{Z}})$ also generates a discrete non-compact subgroup in S_1/S_2 ($\tilde{\rho}$ being an algebraic, hence continuous, map). Hence $s_1^{\mathbb{Z}}$ -orbits in S_1/S_2 are closed so that the space $s_1^{\mathbb{Z}} \backslash S_1/S_2$ is countably separated. By Lemma 9, $s^{\mathbb{Z}}$

acts ergodically on $\Delta \backslash \tilde{H}(\mathbb{R})$. By applying Proposition (2.1.11) of [Z] to the $s_1^{\mathbb{Z}}$ -invariant map $\bar{\phi}^* : \Delta \backslash \tilde{H}(\mathbb{R}) \rightarrow s_1^{\mathbb{Z}} \backslash S_1/S_2$, we deduce that the image of ϕ^* is essentially contained in an orbit of $s_1^{\mathbb{Z}}$ in the quotient group S_1/S_2 . Then, by Lemma 11, it follows that ϕ^* is constant.

Therefore, $\phi^*(h) = p \in G'(k')/J$ is constant, and $\phi(h) = \tilde{\rho}(h)(p)$ is a rational map under the assumption that k' is an archimedean local field.

If k' is non-archimedean, then by Theorem 12, the image $\rho(\Delta)$ is contained in a compact group K which acts smoothly on $G'(k')/J$ so that $K \backslash G'(k')/J$ is countably separated. The group Δ acts ergodically on $\tilde{H}(\mathbb{R})/S$. Hence, by Proposition (2.1.11) [Z], the S -invariant map $\bar{\phi} : \Delta \backslash \tilde{H}(\mathbb{R}) \rightarrow K \backslash G'(k')/J$ is essentially constant, and therefore the image of ϕ is essentially contained in an orbit of K .

Since K is a compact subgroup of the p -adic group $G'(k')$, it has a decreasing sequence of open subgroups $(K_n)_{(n \geq 1)}$ (of finite index in K) such that the intersection $\cap_{n \geq 1} K_n = \{1\}$ is trivial. Then $\Delta_n = \Delta \cap \rho^{-1}(K_n)$ is of finite index in Δ . By Theorem 12 applied to Δ_n , it follows that the image of ϕ is contained in an orbit of K_n for each $n \geq 1$. But since the subgroups K_n 's converge to the identity subgroup, it follows that the image of ϕ is a singleton. That is, ϕ is constant. $\square$

We now mention two consequences of Fubini's Theorem we will need in the proof of Theorem 6.

Notation 1. Suppose that H is a locally compact Hausdorff second countable topological group with a Haar measure μ and assume that (H, μ) is σ -finite. Suppose that (X, ν) is a σ -finite measure space on which H acts such that the action $H \times X \rightarrow X$ - denoted $(h, x) \mapsto hx$ - is measurable and so that for each $h \in H$, the map $x \mapsto hx$ on X preserves the measure class of ν . Let Z be a measure space and let $f : X \rightarrow Z$ be a measurable map. Given a measure space (X, ν) , we will say that a measurable subset $X_1 \subset X$ is **co-null**, if the complement $X \setminus X_1$ has measure zero with respect to ν .

Lemma 13. *Under the preceding notation, suppose that there exists a co-null subset $X_1 \in X$ such that for each $x \in X_1$, the map $h \mapsto f(hx)$ is constant on a co-null subset H_x of H .*

Then, given $h \in H$, there exists a measurable subset X_h which is co-null in X_1 , such that

$$f(hx) = f(x) \quad \forall x \in X_h.$$

Proof. This is essentially a consequence of Fubini's Theorem. Define $E = \{(h, x) \in H \times X : hx \in X_1\}$. Then, for each $h \in H$, we have $E_h \stackrel{\text{def}}{=} \{x \in X : hx \in X_1\} = h^{-1}(X_1)$. Therefore E_h is co-null and hence by Fubini's Theorem, E is co-null in $H \times X$.

Given $h_0 \in H$, $y \in X_1$ and $h \in H$, consider the set

$$W_y = \{h \in H : f(h_0hy) = f(hy)\}.$$

Since $W_y \supset h_0^{-1}H_y \cap H_y$, it follows that W_y is co-null in H . Consider the set $E_1 = \{(h, y) \in E : y \in X_1, \text{ and } f(h_0hy) = f(hy)\}$. Then E_1 is measurable and co-null in E by Fubini, since, for each $y \in X_1$, the set E_1^y is nothing but W_y and is co-null. Write $X_{h_0} \stackrel{\text{def}}{=} \{hy : (h, y) \in E_1\}$. Then, $X_{h_0} \subset X_1$ (since for every $(h, y) \in E$ we have $hy \in X_1$) and for almost all h , we have $(E_1)_h = \{y \in X_1 : hy \in X_{h_0}\} = h^{-1}X_{h_0}$ is measurable and co-null. Hence X_{h_0} is measurable and co-null in X_1 .

Let $x \in X_{h_0}$. Then, by the definition of the set X_{h_0} , there exists $(h, y) \in E_1$ such that $x = hy$. Moreover, $f(h_0x) = f(h_0hy) = f(hy) = f(x)$ (the last but one equation holds because $(h, y) \in E_1$). This proves the Lemma. $\square$

Notation 2. Let G be a locally compact Hausdorff second countable group with a σ -finite Haar measure. Equip $G \times G$ with the product measure $\mu \times \mu$. Let Z be a measure space and $f : G \rightarrow Z$ a measurable map.

The following lemma is again a simple application of Fubini's Theorem.

Lemma 14. *With the preceding notation, suppose that given $g \in G$, there exists a measurable co-null subset $X_g \subset X$, such that*

$$f(gx) = f(x) \quad \forall x \in X_g.$$

Then there exists a measurable co-null subset $Y \subset X$ such that $y \mapsto f(y)$ is constant on Y .

Proof. Let $E = \{(g, x) \in G \times G : f(gx) = f(x)\}$. Then E is measurable and for almost all $g \in G$, the set $E_g = \{x \in G : (g, x) \in E\}$ is measurable. Moreover, by the definition of E , $E_g = \{x \in G : f(gx) = f(x)\}$ and hence contains the co-null subset X_g . Therefore, E_g is co-null for almost all $g \in G$, which implies, by the Fubini Theorem, that E is co-null in $G \times G$. Hence, again by the Fubini Theorem, there exists a co-null subset X_1 in G such that for all $x \in X_1$, we have $E^x = \{g \in G : (g, x) \in E\}$ is co-null in G . But by the definition

of E , $E^x = \{g \in G : f(gx) = f(x)\}$. Fix $x \in X_1$. Set $Y = E^x x$. Then Y is co-null in G . Given $y \in Y$, there exists $g \in E^x$ such that $y = gx$. Therefore, $f(y) = f(gx) = f(x)$ for all $y \in Y$. This proves the lemma. $\square$

Finally, we recall a well known result of Furstenberg we will need, commonly known as Furstenberg's Lemma but due to Zimmer in the form given below.

Lemma 15. (*Furstenberg, see [Z], Corollary (4.3.7) and Proposition (4.3.9)*). *Suppose that Γ is a closed subgroup of a locally compact topological group and that P_0 is a closed amenable subgroup of G . Let X be a compact metric Γ -space. Then there exists a Borel measurable Γ -equivariant map from G/P_0 to $\mathcal{P}(X)$, the space of probability measures on X .*

3. PROOF OF THE SUPER-RIGIDITY RESULT (THEOREM 5)

We will now proceed to the proof of Theorem 5.

In this section, we suppose that H is a semi-simple Lie subgroup of a **simple** Lie group G . Let P be a minimal real parabolic subgroup of G and S and N denote respectively a maximal real split torus of P and the unipotent radical of P . Let A be the connected component of identity in S . Denote by P_0 the subgroup AN of P (if G is split, then $P_0 = P$; in general we have replaced the minimal parabolic subgroup P by a subgroup P_0 which has no compact factors such that P/P_0 is compact). Let K be a maximal compact subgroup of G . We have the Iwasawa decomposition $G = P_0 K = ANK$.

We will treat the archimedean and non-archimedean cases separately.

We start by mentioning the following easy observation.

Lemma 16. *Let H be a semi-simple subgroup of simple group G and K a maximal compact subgroup of G . assume that $\dim(H) > \dim(K)$. Let $G = NAK$ be an Iwasawa decomposition of G and $P_0 = NA$. Then the isotropy subgroup of H at any point in G/P_0 is a non-compact subgroup of H .*

Proof. Since $G/P_0 = K$, we have $\dim(G/P_0) = \dim(K)$, and since $\dim(H) > \dim(K)$, at any point $p \in G/P_0$, the isotropy of H is a positive dimensional subgroup, which is conjugate to a subgroup of P_0 ; the latter has no compact subgroups, hence the isotropy of H at p is a non-compact subgroup. $\square$

Theorem 17. *Suppose Γ is a Zariski dense discrete subgroup of a simple Lie group G which intersects a semi-simple Lie subgroup H (of real rank at least two) of G in an irreducible lattice. Suppose that H acts with non-compact isotropy at any point of G/P_0 (or that $\dim(H) > \dim(K)$ for a maximal compact subgroup K of G). Then, the group Γ is non-archimedean super-rigid.*

Proof. Suppose that $\rho : \Gamma \rightarrow G'$ is a representation of Γ into an absolutely simple algebraic group G' over a **non-archimedean** local field k' of characteristic zero with Zariski dense image.

By a construction of Furstenberg (Lemma 15), there exists a Γ equivariant measurable map ϕ from G/P_0 into $\mathcal{P}(G'(k')/P'(k'))$ where $\mathcal{P}$ refers to the space of probability measures on the relevant space. Being a closed subgroup of P , the group P_0 is amenable.

In particular, the map ϕ is Δ -equivariant. By assumption (or by Lemma 16), at any point $p \in G/P_0$, the isotropy of H is a non-compact subgroup.

Now by Corollary (3.2.17) of [Z], the group $G'(k')$ acts smoothly on $\mathcal{P}(G'(k')/P'(k'))$ and hence the quotient space $\mathcal{P}(G'(k')/P'(k'))/G'(k')$ is countably separated. By Lemma 9, Δ acts ergodically on $\tilde{H}(\mathbb{R})/S$. Therefore, the map

$$\phi_p : H/S \rightarrow \mathcal{P}(G'(k')/P'(k')), h \mapsto \phi(hp)$$

is essentially contained in a $G'(k')$ -orbit (Proposition (2.1.11) of [Z]). Write this orbit as $G'(k')/J$ for some closed subgroup J of $G'(k')$. Then, by Proposition 8, the map $\phi : H/S \rightarrow G'(k')/J$ is constant. Therefore, by Lemma 13, given $h \in H$, there exists a co-null set X_1 in G/P_0 , such that $\phi(hp) = \phi(p)$ for all $p \in X_1$.

Since G is simple and Γ is Zariski dense, there exist finitely many elements $\gamma_1, \gamma_2, \dots, \gamma_k \in \Gamma$ such that the span of the Lie algebras of the conjugates ${}^{\gamma_i}(H) = \gamma_i H \gamma_i^{-1}$ is the Lie algebra of G . The existence of such a set follows from the simplicity of G and the Zariski density of Γ in G . Then, we apply the conclusion of the preceding paragraph successively to the groups ${}^{\gamma_i}(H)$ to conclude that if $h_i \in {}^{\gamma_i}(H)$ ($1 \leq i \leq k$), then the map $x \mapsto \phi(h_1 h_2 \dots h_k x)$ does not depend on the h_i , on a co-null set X_1 . Since the product set ${}^{\gamma_1}(H) {}^{\gamma_2}(H) \dots {}^{\gamma_k}(H)$ contains a Zariski open set in G by the choice of the γ_i , by replacing the set $\{\gamma_i\}$ with a larger set if necessary, we may assume that the product set ${}^{\gamma_1}(H) \dots {}^{\gamma_k}(H) = G$. It follows from Lemma 14 that the map ϕ is

constant, and hence $\rho(\Gamma)$ fixes a probability measure on $G'(k')/P'(k')$.

But the isotropy subgroup in $G'(k')$ of a probability measure μ on $G'(k')/P'(k')$ is (by [Z]) an extension of an algebraic subgroup G'_μ of G' by a compact subgroup of $G'(k')$. Since $\rho(\Gamma)$ is Zariski dense in $G'(k')$ it follows that G'_μ is normalised by G' ; since G' is absolutely simple, it follows that G'_μ is trivial, and $\rho(\Gamma)$ is contained in a compact subgroup of $G'(k')$.

This means that Γ is non-archimedean super-rigid in G , and proves Theorem 17. $\square$

To treat the non-archimedean case, we need the following preliminary result.

Lemma 18. *If $f : X \rightarrow Y$ is a surjective map of irreducible affine varieties over $\mathbb{R}$, such that $X(\mathbb{R})$ (resp. $Y(\mathbb{R})$) is Zariski dense in X (resp. Y), and $\phi : Y(\mathbb{R}) \rightarrow \mathbb{R}$ is a set theoretic map such that the composite $\phi \circ f$ is a rational map on $X(\mathbb{R})$, then ϕ is a rational map on $Y(\mathbb{R})$.*

Proof. We may assume - by replacing X by an open set - that X is a finite cover $\tilde{\mathcal{U}}$ of an open set $\mathcal{U}$ of a product $Y \times Z$ of Y with an affine variety Z ; and that the map f is the composite of the covering map $p : \tilde{\mathcal{U}} \rightarrow \mathcal{U}$ with the first projection $\pi_1 : Y \times Z \rightarrow Y$.

The assumptions imply that the composite $p \circ \pi_1 \circ \phi$ is rational on $\tilde{\mathcal{U}}$. Since p is a finite cover, it follows that $\pi_1 \circ \phi$ is rational on $\mathcal{U}$ (a rational map on a finite cover which descends to a function on the base is a rational function on the base), and hence rational on $Y \times Z$. Since this map is independent of the point $z \in Z$, it follows that the map ϕ is rational on Y (a rational function in two variables which depends only on the first variable is a rational function in the first variable). $\square$

Theorem 19. *Let H be a semi-simple Lie subgroup (of real rank at least two) of a simple Lie group G which acts with non-compact isotropies on G/P_0 (or, which satisfies the stronger condition $\dim(H) > \dim(K)$ for a maximal compact subgroup K of G). Let $\Gamma \subset G$ be a Zariski dense discrete subgroup which intersects H in an irreducible lattice. Let $\rho : \Gamma \rightarrow G'(k')$ be a homomorphism of Γ , with k' an archimedean local field, and G' an absolutely simple algebraic group over k' . If $\rho(\Gamma)$ is not relatively compact in $G'(k')$ and is Zariski dense in G' , then ρ*

extends to an algebraic homomorphism of G into $G'(k')$ defined over $\mathbb{R}$.

Proof. If ρ is an archimedean representation of Γ we first prove that on each H -orbit, the map ϕ is a rational map. More precisely, we prove that the map ϕ is the composite of an algebraic homomorphism of $\tilde{H}$ into $G'(k')$ with the orbit map $G'(k') \rightarrow G'(k')p$ with $p \in \mathcal{P}$. This is exactly the statement of Proposition 8 if $H = \tilde{H}(\mathbb{R})$, where $\tilde{H}$ is simply connected. If not, we replace H (resp. Δ) by the image of $\tilde{H}(\mathbb{R})$ where $\tilde{H}$ is the simply connected cover of H (resp. $\Delta \cap \tilde{H}(\mathbb{R})$). The conclusion of Proposition 8 is unaltered.

As in the proof of Theorem 17, we fix elements $\gamma_1, \gamma_2, \dots, \gamma_k \in \Gamma$ such that the Lie algebras of the conjugates ${}^{\gamma_i}(H) = \gamma_i H \gamma_i^{-1}$ together span that of G . The existence of such a set follows from the simplicity of G and the Zariski density of Γ in G . Then, we apply the conclusion of the preceding paragraph successively to ${}^{\gamma_i}(H)$ to conclude that if $h_i \in {}^{\gamma_i}(H)$ (with $1 \leq i \leq k$), then the map $x \mapsto \phi(h_1 h_2 \cdots h_k x)$ is simply the map

$$\tilde{\rho}(h_1) \tilde{\rho}(h_2) \cdots \tilde{\rho}(h_k) (\phi(x)),$$

for almost all $x \in G/P_0$, and all $h_i \in H_i$. Here

$$\tilde{\rho}(h_i) = \rho(\gamma_i) \tilde{\rho}(\gamma_i^{-1} h_i \gamma_i) \rho(\gamma_i)^{-1}$$

is the natural homomorphism on the conjugate ${}^{\gamma_i}(H)$ into $G'(k')$ obtained from $\tilde{\rho} : H \rightarrow G'(k')$.

Since the product set ${}^{\gamma_1}(H) {}^{\gamma_2}(H) \cdots {}^{\gamma_k}(H)$ contains a Zariski open set in G by the choice of the γ_i , by replacing the product by a product over a larger set of γ_i 's if necessary, we may assume that the product set ${}^{\gamma_1}(H) {}^{\gamma_2}(H) \cdots {}^{\gamma_k}(H)$ contains all of G (we may assume that G is connected). Given $g \in G$, there exist elements $h_i \in {}^{\gamma_i}(H)$ ($1 \leq i \leq k$) such that $g = h_1 \cdots h_k$. Thus, given $g \in G$, there exists an element $R(g) = \tilde{\rho}(h_1) \cdots \tilde{\rho}(h_k) \in G'(k')$ such that for a co-null set $X_1 \subset G/P_0$, we have

$$(1) \quad \phi(gx) = R(g) \phi(x) \quad \forall x \in X_1.$$

We may assume (since Γ is countable), that X_1 is Γ -stable. If $R'(g) \in G'(k')$ is another element such that the equality (1) holds with $R'(g)$ in place of $R(g)$, then, we have: $R(g)^{-1} R'(g) \in I_{\phi(x)}$, the isotropy of $G'(k')$ at the point $\phi(x)$, for each $x \in X_1$. Replacing x by γx for a fixed $\gamma \in \Gamma$, $R(g)^{-1} R'(g)$ lies in the intersection

$$I = \bigcap_{\gamma \in \Gamma} \rho(\gamma) I_{\phi(x)} \rho(\gamma)^{-1}.$$

But the isotropy is an algebraic subgroup of $G'(k')$ (by [Z]). Hence the intersection I is also algebraic, but is normalised by the Zariski dense subgroup $\rho(\Gamma)$ of $G'(k')$, and hence is normalised by $G'(k')$. But G' is absolutely simple, hence the intersection is trivial and $R(g) = R'(g)$ for all $g \in G$. In particular, $R(\gamma) = \rho(\gamma)$ for all $\gamma \in \Gamma$.

Moreover, the equation

$$\phi(g_1 g_2 x) = R(g_1) R(g_2) \phi(x) = R(g_1 g_2) \phi(x)$$

holds for a fixed $g_1, g_2 \in G$, and $x \in X'$ a co-null subset of G/P_0 . By the uniqueness of $R(g_1 g_2)$ proved in the preceding paragraph, we have: $R(g_1 g_2) = R(g_1) R(g_2)$ for any fixed pair $g_1, g_2 \in G$. Hence $R : G \rightarrow G'(k')$ is an abstract homomorphism.

The composite of the product map $f : H_1 \times \cdots \times H_k \rightarrow G$ and the abstract homomorphism $R : G \rightarrow G'(k')$ is $R \circ f(h_1, \dots, h_k) = \tilde{\rho}(h_1) \cdots \tilde{\rho}(h_k)$, which is rational on $H_1 \times \cdots \times H_k$. Then the map R is rational, by virtue of Lemma 18.

Theorem 19 now follows since the rationality of R implies the rationality of the representation $\rho : \Gamma \rightarrow G'(k')$. $\square$

Theorem 5 is an immediate consequence of Theorem 17 and Theorem 19. Theorem 6 is a particular case of Theorem 5, in view of Lemma 16.

4. APPLICATIONS (PROOF OF COROLLARY 1)

Assume that $H = SL_k(\mathbb{R})$ and $G = SL_n(\mathbb{R})$. Under the assumptions of Corollary 1, we have $k - 1 > n/2$ and $\dim(H) = k^2 - 1 > \dim(G/P_0) = n(n - 1)/2$. Therefore, if Γ is a Zariski dense discrete subgroup of G which intersects H in a lattice, then by Theorem 6, Γ is super-rigid. We now prove Corollary 1.

We now recall a result, which is a generalisation of Margulis' observation that super-rigidity implies arithmeticity. However, Margulis needed only that the discrete subgroup was a lattice. We have not assumed that Γ is a lattice (indeed, this is what is to be proved), and we also do not assume that Γ is finitely generated.

Theorem 20. ([V3]) *Let G be an absolutely simple real algebraic group and let Γ be a super-rigid discrete subgroup. Then there exists an arithmetic group Γ_0 of G containing Γ .*

Suppose $\Gamma_0 \subset G$ is arithmetic. This means that there exists a number field F and a semi-simple linear algebraic F -group $\mathbf{G}$ such that the

group $\mathbf{G}(\mathbb{R} \otimes F)$ is isomorphic to a product $SL_n(\mathbb{R}) \times U$ of $SL_n(\mathbb{R})$ with a compact group U . Under this isomorphism, the projection of $\mathbf{G}(O_F)$ of the integral points of $\mathbf{G}$ into G is commensurable with Γ_0 . The simplicity of $SL_n(\mathbb{R})$ implies that $\mathbf{G}$ may be assumed to be absolutely simple over F . The group $\mathbf{G}$ is said to be an F -form of SL_n . Moreover, if Γ_0 contains unipotent elements, then $\mathbf{G}$ can not be anisotropic over F . Hence F -rank of $\mathbf{G}$ is greater than zero. In that case, $\mathbf{G}(F \otimes \mathbb{R})$ can not contain compact factors (since compact groups can not contain unipotent elements). This means that $F = \mathbb{Q}$.

We now recall the classification of $\mathbb{Q}$ -forms of SL_n .

[1] Let d be a divisor of n and D a central division algebra over $\mathbb{Q}$ of degree d . Write $n = md$. Then, the algebraic group $\mathbf{G} = SL_m(D)$ is a $\mathbb{Q}$ -form of SL_n . The rank of $\mathbf{G}$ is $m - 1 = n/d - 1$. If $d \geq 2$, then $m - 1 < n/2$.

[2] Let $E/\mathbb{Q}$ be a quadratic extension and D a central division algebra over E with an involution of the second kind with respect to $E/\mathbb{Q}$. Let d be the degree of D over E , suppose d divides n and let $md = n$. Let $h : D^m \times D^m \rightarrow E$ be a Hermitian form with respect to this involution, and let $\mathbf{G} = SU(h)$. Then, $\mathbf{G}$ is a $\mathbb{Q}$ -form of SL_n ; its $\mathbb{Q}$ -rank is not more than $m/2 = n/2d \leq n/2$.

The classification of simple algebraic groups (see [T]), implies the following.

Lemma 21. *The only $\mathbb{Q}$ -forms $\mathbf{G}$ of SL_n are as above. In particular, if $\mathbf{G}$ is a $\mathbb{Q}$ -form of $\mathbb{Q}$ -rank strictly greater than $n/2$, then $\mathbf{G}$ is $\mathbb{Q}$ -isomorphic to SL_n .*

Proof of Corollary 1. By Theorem 6, the group Γ is superrigid in G . By Theorem 20, Γ is contained in an arithmetic subgroup Γ_0 of G . Since $\Gamma_0 \supset \Gamma$ contains a finite index subgroup of $SL_k(\mathbb{Z})$ by assumption, it follows that Γ_0 contains unipotent elements. Therefore, the number field F associated to Γ_0 is $\mathbb{Q}$. Since H intersects Γ_0 in $SL_k(\mathbb{Z})$, it follows that H is the group of real points of a $\mathbb{Q}$ -subgroup $\mathbf{H}$ of $\mathbf{G}$. Clearly $\mathbf{H} = SL_k$ since it contains $SL_k(\mathbb{Z})$ in its $\mathbb{Q}$ -rational points. The $\mathbb{Q}$ -rank of $\mathbf{G}$ is not less than the $\mathbb{Q}$ -rank of $\mathbf{H}$ which is $k - 1 > n/2$ by assumption.

By Lemma 21, the $\mathbb{Q}$ -form $\mathbf{G}$ is isomorphic to SL_n . Hence Γ_0 is commensurable with $SL_n(\mathbb{Z})$. Moreover, the $\mathbb{Q}$ -inclusion of $\mathbf{H} = SL_k$

in $\mathbf{G} = SL_n$ is the standard one described before the statement of Theorem 3.

Now, Γ is Zariski dense and contains $SL_k(\mathbb{Z})$. Let $e_1, e_2, \dots, e_n$ be the standard basis of $\mathbb{Q}^n$. Consider the change of basis which interchanges e_k and e_n and all other e_i 's are left unchanged. After this change of basis, (which leaves the diagonal torus stable), the group Γ (or rather, a conjugate of it by the matrix effecting this change of basis) contains the highest root group and the second highest root group (in the usual notation for SL_n the positive roots occur in the Lie algebra of upper triangular matrices). By Theorem (3.5) (or Corollary (3.6)) of [V1], Γ must be of finite index in $SL_n(\mathbb{Z})$. This proves Corollary 1. $\square$

Corollary 2 is proved in an analogous way.

5. PROOF OF THEOREM 3

Notation 3. Let $k \geq 3$ and $m \geq 2$ be integers, set $n = k + m$. Fix (real) vector spaces W , E and V of dimensions k , m and n respectively. Assume that $V = W \oplus E$. Then there is a decomposition of the dual vector spaces $V^* = W^* \oplus E^*$. Fix a basis $w_1, w_2, \dots, w_k$ of the vector space W . Let $w_1^*, w_2^*, \dots, w_k^*$ be the dual basis in $W^* \subset V^*$. Let

$$\mathcal{U} = \{g \in SL(V) : g(w_i) \notin W (1 \leq i \leq k-1) \text{ and } g^*(w_k^*) \notin E^* \cup W^*\}.$$

Then, $\mathcal{U}$ is a non-empty set in G which is open in the Zariski topology on $SL(V)$.

We will view $SL(W)$ as a subgroup of $SL(V)$ by letting any element of $SL(W)$ act trivially on E . Let $U \subset SL(W)$ be the unipotent subgroup an element of which acts trivially on each w_i with $1 \leq i \leq k-1$ and takes w_k to w_k plus a multiple of w_1 . The subspace of vectors in V fixed by U . Denote by $L(g)$ the Zariski closure in $SL(V)$ of the subgroup generated by $SL(W)$ and gUg^{-1} .

Lemma 22. *With the above notation, (and the assumption that $k \geq 3$), there exists a Zariski open set $\mathcal{U}$ in $SL_n(\mathbb{R})$ such that the quotient of the group $L(g)$ modulo its radical is isomorphic to SL_{k+1} , for every $g \in \mathcal{U}$.*

Proof. The lemma is equivalent to showing that the Lie algebra $l(g)$ generated by $sl(W)$ and $g(lie(U))g^{-1}$ in the Lie algebra $sl(V)$ of the group $SL(V)$. We first decompose the space $sl(V)$ as a module over

$sl(W)$. Now, $V = W \oplus \mathbb{C}^{n-k}$ where $\mathbb{C}$ is the one dimensional trivial $sl(W)$ -module. Hence

$$sl(V) = sl(W) \oplus \mathbb{C}^{n-k} \otimes W \oplus W^* \otimes \mathbb{C}^{n-k} \oplus E$$

where E is a trivial $sl(W)$ module of the appropriate dimension and W^* is the dual of W .

The Lie algebra $Lie(U)$ is generated by the linear transformation E_{k1} which takes all the elements of the standard basis of $\mathbb{R}^n$ to zero, except the k -th basis element w_k which is taken to w_1 . Therefore, the kernel of E_{k1} has codimension 1. By the genericity assumption on g , the conjugate $gE_{k1}g^{-1}$ has kernel $F(g)$ (also of codimension 1) which intersects E in a subspace of codimension 1: $\dim(F(g) \cap E) = n - k - 1$.

We have the exact sequence

$$0 \rightarrow F(g) \cap E \rightarrow V \rightarrow V/F(g) \cap E \rightarrow 0$$

of vector spaces, which is also a sequence of $l(g)$ modules since all the terms are stable under the action of $sl(W)$ and $gLie(U)g^{-1}$. Moreover, $l(g)$ acts trivially on $F(g) \cap E$. Denote by $\overline{V}$ the quotient $V/F(g) \cap E$, by $\overline{E}$ the image of $E \subset V$ in $\overline{V}$. Then $\overline{E}$ is one dimensional.

Moreover, as an $sl(W)$ -module, we have the decomposition $\overline{V} = W \oplus \overline{E}$ and hence as an $sl(W)$ -module, the decomposition

$$End(\overline{V}) = End(W) \oplus \overline{E} \otimes E \oplus W^* \otimes \overline{E} \oplus \overline{E}^* \otimes W.$$

Since $\dim(W) \geq 3$, the irreducible modules W and W^* are non-isomorphic; therefore, if a vector in $sl(\overline{V})$ projects non-trivially to $W \otimes \overline{E}^*$ and to $\overline{E} \otimes W^*$, then the $sl(W)$ -module generated by the vector contains both the spaces $W \otimes \overline{E}^*$ and $\overline{E} \otimes W^*$.

The genericity assumption on g ensures that the projection of $gE_{k1}g^{-1}$ viewed as an endomorphism of $\overline{V}$ projects non-trivially to W and W^* in $End(\overline{V})$. Hence, by the preceding paragraph, $l(g)$ contains in its image in $End(\overline{V})$, $sl(W)$, W^* and W ; therefore, the image of $l(g)$ equals $sl(\overline{V})$. This proves the Lemma. $\square$

Lemma 23. *If $\Delta \subset \mathcal{G} = \mathbb{R}^{k+1} \times SL_{k+1}(\mathbb{R})$ is a Zariski dense discrete subgroup of $\mathcal{G}$ which contains $\{0\} \times SL_{k-1}(\mathbb{Z})$ embedded in $SL_k(\mathbb{R})$ in the top left hand corner, and if $k \geq 3$, then, the group Δ virtually contains a conjugate of $SL_{k+1}(\mathbb{Z})$.*

Proof. The group Δ may be shown to be super-rigid in $\mathcal{G}$, exactly as in the proof of Corollary 1, since $k \geq 3$. This means, by [V3], that the projection of Δ is contained in an arithmetic subgroup of $SL_{k+1}(\mathbb{R})$, and is *therefore* discrete. The Zariski density of the projection in $SL_{k+1}(\mathbb{R})$ now implies, by Corollary 1, that the *projection* of Δ is conjugate virtually to $SL_{k+1}(\mathbb{Z})$.

Now, Δ is a subgroup of a semi-direct product and its projection to $SL_{k+1}(\mathbb{R})$ contains virtually $SL_{k+1}(\mathbb{Z})$. Hence Δ defines a element of the cohomology group $H^1(\Delta', \mathbb{R}^n)$ where $\Delta' \subset SL_{k+1}(\mathbb{Z})$ is a subgroup of finite index. But the first cohomology group vanishes, by a Theorem of Raghunathan ([R2]), for $k + 1 \geq 3$, and therefore, Δ contains, virtually, a conjugate of $SL_{k+1}(\mathbb{Z})$. $\square$

We will now begin the proof of Theorem 3. We have $SL_3(\mathbb{Z})$ is virtually contained (in the top left hand corner) in the Zariski dense discrete subgroup Γ of $SL_n(\mathbb{R})$. We will prove by induction, that for **every** $k \geq 3$ with $k \leq n$, a conjugate of $SL_k(\mathbb{Z})$ is virtually a subgroup of Γ . Applying this to $k = n$, we get Theorem 3.

Suppose that for some k , we have $SL_k(\mathbb{Z})$ is virtually contained in Γ . Let $\gamma \in \Gamma$, and let $\Delta = \Delta(\gamma)$ denote the subgroup of Γ generated by $SL_k(\mathbb{Z})$ and a conjugate of the unipotent element $\gamma u \gamma^{-1}$ (where u is in a finite index subgroup of $SL_3(\mathbb{Z})$, as in Lemma 22). Assume further, that the element γ is in the open set $\mathcal{U}$ of Notation 3. By Lemma 22, the Zariski closure of the group Δ maps onto SL_{k+1} .

By Lemma 23, the group Δ contains, virtually, the group $SL_{k+1}(\mathbb{Z})$. We have thus proved that if $SL_k(\mathbb{Z}) \subset \Gamma$, then $SL_{k+1}(\mathbb{Z}) \subset \Gamma$, provided $k + 1 \leq n$. Thus the induction is completed and therefore Theorem 3 is established

6. THE RANK ONE CASE

In this last section, we will see that the situation for Nori's question is completely different in real rank one. More precisely, we have the following result.

Theorem 24. *Let G be a real simple Lie group of real rank one and $H \subset G$ a non-compact semi-simple subgroup. Suppose that Δ is a lattice in H . Then, there exists a Zariski dense discrete subgroup Γ in G of infinite co-volume whose intersection with H is a subgroup of finite index in the lattice Δ .*

Suppose that H is a simple non-compact subgroup of a simple group G of real rank one. Let P be a minimal parabolic subgroup of G which intersects H in a minimal parabolic subgroup Q . The group G acts on G/P and H leaves the open set $\mathcal{U} = G/P \setminus H/Q$ stable. Let Δ be a discrete subgroup of H .

Lemma 25. *Given compact subsets Ω_1 and Ω_2 in the open set $\mathcal{U}$, the set $\mathcal{U}_H = \{h \in H : h\Omega_1 \subset \Omega_2\}$ is a compact subset of H .*

Further, the group Δ acts properly discontinuously on $\mathcal{U}$.

Proof. Choose, as one may, a maximal compact subgroup K of G such that $K \cap H$ is maximal compact in H . The set $\mathcal{U}$ is invariant under H and hence under $K \cap H$. We may assume that the compact sets Ω_1 and Ω_2 are invariant under $K \cap H$ of H . The Cartan decomposition of H says that we may write $H = (K \cap H)A^+(K \cap H)$ with A a maximal real split torus (of dimension one) and $A^+ = \{a \in A : 0 < \alpha(h) \leq 1\}$ where α is a positive root of A . Let W denote the normaliser modulo the centraliser of A in G . This is the relative Weyl group. Since $\mathbb{R}$ -rank $(G)=1$, it follows that $W = \{1, \kappa\}$ has only two elements. Since $\mathbb{R}$ -rank $(H)=1$, it follows that $\kappa \in H$. We have the Bruhat decomposition $G = P \cup U\kappa P$.

If possible, let h_m be a sequence in $\mathcal{U}_H$ which tends to infinity. It follows from the previous paragraph that $h_m = k_m a_m k'_m$ with $k_m, k'_m \in K \cap H$ and $a_m \in A^+$, and $\alpha(a_m) \rightarrow 0$ as $m \rightarrow +\infty$. Let p be an element of $\Omega_1 \subset \mathcal{U}$. By the Bruhat decomposition in G (G has real rank one), it follows that $\mathcal{U} \subset U\kappa P$, where κ is a non-trivial element of the Weyl group of A in G and U is the unipotent radical of the minimal parabolic subgroup P . We may write $p = ukP$ for some $u \in U$. Since $h_m p \in \Omega_2$ and the latter is compact, we may replace h_m by a sub-sequence and assume that $h_m p$ converges, say to $q \in \Omega_2$. Since Ω_2 is stable under $K \cap H$, we may assume that $k_m = 1$, by replacing q by a sub-sequential limit of $k_m^{-1}q$.

We compute $h_m p = a_m k'_m u \kappa P \in \Omega_2 \subset \mathcal{U}$. The convergence of k'_m says that $k'_m u \kappa P = u_m \kappa P$ with u_m convergent (possibly after passing to a sub-sequence). We write ${}^a(u) \stackrel{\text{defn}}{=} aua^{-1}$. Then, $h_m p = {}^{a_m}(u_m) \kappa P$ since κ conjugates A into P (in fact into A). Since $\alpha(a_m) \rightarrow 0$, and u_m are bounded, conjugation by a_m contracts u_m into the identity and $h_m p$ therefore tends to κP . But the latter is in H/Q since κ already lies in H . Hence the limit does not lie in Ω_2 , contradicting the fact that Ω_2 is compact. Therefore $\mathcal{U}_H$ is compact.

The second part of the lemma immediately follows since the intersection of Δ with $\mathcal{U}_H$ is finite. $\square$

Remark 1. Observe that if g does not lie in the product set HP (which is a Zariski closed subset of measure zero in G (since $HP/P = H/Q$ is closed and of measure zero in G/P), then $g(HP/P) \cap HP/P = \emptyset$. Hence $g(HP/P)$ is a compact subset of $\mathcal{U}$ if $g \notin HP$. Now $G \setminus HP$ is a Zariski open subset of G and hence intersects any Zariski dense subset non-trivially.

Given a lattice Δ in H , and given a compact subset $\Omega \subset \mathcal{U}$, Lemma 25 shows that there exists a finite index subgroup Δ' of Δ such that non-trivial elements of Δ' drag Ω into an arbitrarily small compact neighbourhood $\mathcal{V}$ of $H/Q = HP/P$.

By Remark 1 there exists $g \in G - HP$ such that $g(\mathcal{V}) \subset \mathcal{U}$. Replacing H/Q by $g(H/Q)$ and Δ by $g\Delta g^{-1}$, we see that all points of H/Q are dragged, by non-trivial elements of $g\Delta g^{-1}$, into a small neighbourhood of H/Q . The ping-pong lemma then guarantees that there exists a finite index subgroup Δ'' such that the group Γ generated by Δ'' and $g\Delta''g^{-1}$ is the free product of Δ'' and $g\Delta''g^{-1}$.

We may replace g by a finite set $g_1, \dots, g_k$ such that for each pair i, j , the intersections $g_i(H/Q) \cap g_j(H/Q)$ and $g_i(H/Q) \cap H/Q$ are all empty. By arguments similar to the preceding paragraph, we can find a finite index subgroup Δ_0 of Δ such that the group Γ generated by $g_i\Delta_0g_i^{-1}$ is the free product of the groups $g_i\Delta_0g_i^{-1}$, and by choosing the g_i suitably, we ensure that Γ is Zariski dense in G . This proves Theorem 1, since Γ is discrete and since it operates properly discontinuously on some open set in G/P , Γ can not be a lattice. This proves Theorem 24.

Acknowledgments:

T.N.Venkataramana extends to Madhav Nori his hearty thanks for many very helpful discussions over the years; the present paper addresses a very special case of a general question raised by him about discrete subgroups containing higher rank lattices. He also thanks Marc Burger for explaining the structure of isotropy subgroups of measures on generalized Grassmanians. The excellent hospitality of the Forschungsinstitut für Mathematik, ETH, Zurich while this paper was written is gratefully acknowledged.

The paper was completed during Indira Chatterji's visit to the Tata Institute and she thanks Tata Institute for providing fantastic working conditions.

REFERENCES

- [B] H. Bass, *Representation rigid subgroups*, Pacific Journal on Math. **86** (1980), 15-51.
- [B-L] H. Bass and A. Lubotzky, *Nonarithmetic superrigid groups: counterexamples to Platonov's conjecture*, Ann. of Math.(2) **151** (2000), **3**, 1151-1173.
- [BMS] H. Bass, J. Milnor and J.-P. Serre, *Solution of the congruence subgroup problem for $SL_n(n \geq 3)$ and $Sp_{2n}(n \geq 2)$* , Publ. Math. I.H.E.S., **33** (1967), 59-137.
- [F-K] R. Fricke and F. Klein, *Vorlesungen Über die theorie der automorphische funktionen*, Johnson Reprint Corp., New York; B.G.Teubner Verlagsgesellschaft, Stuttgart 1965 Band I und II.
- [M] G. A. Margulis, *Discrete subgroups of semi-simple lie groups*, Berlin-Heidelberg-New York, Springer-Verlag, (1991).
- [O] Hee Oh, *Discrete groups generated by lattices in opposite horospherical subgroups*, J. Algebra **203** (1998), **no. 2**, 621-676.
- [P-R] V. P. Platonov and A. Rapinchuk, *Algebraic Groups and Number Theory*, Academic Press, (1994).
- [R] M. S. Raghunathan, *A Note on Generators for arithmetic subgroups of algebraic groups* Pac J. Math. **152** (1992), 365-373.
- [R2] M.S.Raghunathan, *Cohomology of Arithmetic Subgroups of Algebraic Groups (I) and (II)*. Ann of Math, **86**, (1967) 409-424, **87**, (1967), 279-304.
- [T] J. Tits, *Classification of Algebraic Semi-simple Groups*, Algebraic Groups and Discontinuous Groups, Proc. Symp. Pure Math., Amer. Math. Soc. Providence, Rhode Island, Boulder, Colorado, (1965), 33-62.
- [V1] T. N. Venkataramana, *Zariski Dense subgroups of arithmetic groups*, J. of Algebra **108**, (1987), 325-339.
- [V2] T. N. Venkataramana, *On some rigid subgroups of semi-simple Lie groups*, Israel J. Math., **89** (1995), **No. 1-3**, 227-236.
- [V3] T. N. Venkataramana, *On the arithmeticity of certain rigid subgroups* CR.Acad. Aci. Paris. s'er I, Math **316** (1993), **no. 4**, 321-326.

- [Z] R. Zimmer, *Ergodic Theory and Semi-simple groups*, Boston-Basel-Stuttgart, Birkhauser, (1984).

E-mail address: `indira.chatterji@math.cnrs.fr`, `venky@math.tifr.res.in`